

CHAPTER VI

REFERENCES

- Antonarakis, S.E.**, Irkin, S.H., Cheng, T.C., Scott, A.F., Sexton, J.P., Trusko, S.P., Charache, S. & Kazazian, H.H., Jr. (1984). β -thalassemia in American Blacks: novel mutations in the "TATA" box and an acceptor splice site. *Proc Natl Acad Sci U S A*, 81, 1154-8.
- Ballas, S.K.**, Cai, S.P., Gabuzda, T. & Chehab, F.F. (1997). Molecular basis of asymptomatic β -thalassemia major in an African American individual. *Am J Med Genet*, 69, 196-9.
- Breathnach, R.** (1981). Organization and expression of eukaryotic split genes coding for proteins. *Annu Rev Biochem*, 50, 349-383.
- Camaschella, C.** & Cappellini, M.D. (1995). Thalassemia intermedia. *Haematologica*, 80, 58-68.
- Camaschella, C.**, Mazza, U., Roetto, A., Gottardi, E., Parziale, A., Travi, M., Fattore, S., Bacchiega, D., Fiorelli, G. & Cappellini, M.D. (1995). Genetic interactions in thalassemia intermedia: analysis of β -mutations, α -genotype, γ -promoters, and β -LCR hypersensitive sites 2 and 4 in Italian patients. *Am J Hematol*, 48, 82-7.
- Chen, J.**, Liu, W. & Chen, M. (1997). Molecular diagnosis of β -thalassemia intermedia. *Zhonghua Yi Xue Za Zhi*, 77, 575-8.

Chen, M., Han, J.Y., Sun, Q. & et al. (2000). Molecular diagnosis in a Korean family with thalassemia intermedia due to co-inheritance of triplicated α -globin genes [$\alpha\alpha/\alpha\alpha\alpha$ (anti 3.7)] and β -thalassemia trait (IVS- II-1; G-A). *Zhonghua Xue Ye Xue Za Zhi*, 21, 195-7.

Cheng T.C., Orkin, S.H., Antonarakis, S.E. & et al. (1984). β -thalassemia in chinese. Use of in vivo RNA analysis and oligonucleotide hybridization in systematic characterization of molecular defects. *Proc Natl Acad Sci U S A*, 81, 2821.

Chernoff, A.I., Minnich, V. & Chongchareonsuk, S. (1954). Hemoglobin E, a hereditary abnormality of human hemoglobin. *Science*, 120, 605.

Clegg, JB. (1987). Can the product of the ζ -gene be a real globin?. *Nature*, 329, 465.

Bio-Rad Laboratories, D.G. (1994). Variant TM β -thal Short program.

Eckman, J. (1998). Common hemoglobin variants facts. *Genetic Northwest*, 7, 2.

Embry SH., Miller, JA. & Dozy AM. (1980). Two different molecular organizations account for the single α -globin gene of the α -thalassemia-2 genotype. *J Clin Invest*, 66, 1319-1325.

Fischel-Ghodsian, N., Vickers, M.A., Seip M., Winichagoon, P. & Higgs, D.R. (1988). Characterization of two deletions that remove the entire human ζ - α globin gene complex. *Br J Haematol*, 70.

Fonseca, S.F., Kerbauy, J., Escrivao, C., Figueiredo, M.S., Cancado, R., Arruda, V.R., Saad, S.T. & Costa, F.F. (1998). Genetic analysis of β -thalassemia major and β -thalassemia intermedia in Brazil. *Hemoglobin*, 22, 197-207.

Fucharoen, S., Fucharoen, G., Sriroongrueng, W., Laosombat, V., Jetsrisuparb, A., Prasatkaew, S., Tanphaichitr, V.S., Suvatte, V., Tuchinda, S. & Fukumaki, Y. (1989). Molecular basis of β -thalassemia in Thailand: analysis of β -thalassemia mutations using the polymerase chain reaction. *Hum Genet*, 84, 41-6.

Fucharoen, S., Ketvichit, P., Pootrakul, P., Siritanaratkul, N., Piankijagum, A. & Wasi, P. (2000). Clinical manifestation of β -thalassemia/hemoglobin E disease. *J Pediatr Hematol Oncol*, 22, 552-7.

Fucharoen, S. & Winichagoon, P. (2002). Thalassemia and abnormal hemoglobin. *Haematology*, supplement II, 83-89.

Fuebetta, M., Tuveri, T., Rosatacelli, C. & et al. (1983). Molecular accounting for milder types of thalassemia major. *J Pediatr*, 103, 35.

Galanello, R. & Cao, A. (1998). Relationship between genotype and phenotype. Thalassemia intermedia. *Ann N Y Acad Sci*, 850, 325-33.

Galanello, R., Dessi, E., Melis, M.A., Addis, M., Sanna, M.A., Rosatelli, C., Argioli, F., Giagu, N., Turco, M.P., Cacace, E. & et al. (1989). Molecular analysis of β^0 -thalassemia intermedia in Sardinia. *Blood*, 74, 823-7.

Garewal, G., Fearon, C.W., Warren, T.C., Marwaha, N., Marwaha, R.K., Mahadik, C. & Kazazian, H.H., Jr. (1994). The molecular basis of β -thalassaemia in Punjabi and Maharashtran Indians includes a multilocus aetiology involving triplicated α -globin loci. *Br J Haematol*, 86, 372-6.

Gilman, J.G. & Huisman, T.H.J., (1984). Two independent genetic factors in the β -globin gene cluster are associated with high $^G\gamma$ -levels in the HbF of SS patients. *Blood*, 64, 452.

Gregory, R.C., Taxman, D.J., Seshasayee, D., Kensinger, M.H., Bieker, J.J. & Wojchowski, D.M. (1996). Functional interaction of GATA1 with erythroid Kruppel-like factor and Sp1 at defined erythroid promoters. *Blood*, 87, 1793.

Ho, P.J., Hall, G.W., Luo, L.Y., Weatherall, D.J. & Thein, S.L. (1998). Beta-thalassaemia intermedia: is it possible consistently to predict phenotype from genotype? *Br J Haematol*, 100, 70-8.

Jandl, J.W. (1987). Method for Reticulocyte Counting. Proposed standard H16-P. *Blood*, 51, 33.

Kazazian H.H., Jr., Orkin, S.H., Antonarakis, S.E., Sexton, J.P., Boehm, C.D., Goff, S.C., Waber, P.G. (1984). Molecular characterization of seven β -thalassemia mutation in Asian Indians. *EMBO J*, 3, 393-396.

Kulozik, A.E., Kohne, E. & Kleihauer, E. (1993). Thalassemia intermedia: compound heterozygous β^0/β^+ -thalassemia and co-inherited heterozygous α^+ -thalassemia. *Ann Hematol*, 66, 51-4.

Kutlar, A., Kutlar, F., Gu, L.G., Mayson, S.M., Huisman, T.H. (1990). Fetal hemoglobin in normal adults and β -thalassemia heterozygotes. *Hum Genet*, 85, 106.

Labie, D., Pagnier, J., Lapoumeroulie, C., et al. (1985). Common haplotype dependency of high γ^G -globin gene expression and high HbF levels in β -thalassemia and sickle cell anemia patients. *Proc Natl Acad Sci U S A*, 82, 2111.

Laig, M., Sanguansermsri, T., Wiangnon, S., Hundrieser, J., Pape, M., Flatz, G. (1989). The spectrum of β -thalassemia mutation in northern and north eastern Thailand. *Hum Genet*, 87, 47-50.

Laosombat, V., Wongchanchailert, M., Sattayasevana, B., Wiriyasateinkul, A. & Fucharoen, S. (2001). Clinical and hematological features of codon 17, A-T mutation of β -thalassemia in Thai patients. *Eur J Haematol*, 66, 126-9.

Lauer J., Shen, C.K.J., Maniatis, T. (1980). The chromosomal arrangement of human α - β -like globin gene. *Cell*, 18, 119-130.

Lawn R.M., Efstratiadis, A., O'Connell, C., Maniatis, T. (1980). The nucleotide sequence of the human-globin gene. *Cell*, 21, 647-651.

Leoni, G.B., Rosatelli, C., Vitucci, A., Addis, M., Loi, A., Tannoia, N. & Cao, A. (1991). Molecular basis of β -thalassemia intermedia in a southern Italian region (Puglia). *Acta Haematol*, 86, 174-8.

Malthis D.J., & Chambon, P. (1981). The SV40 early region TATA box is required for Maltoth Y., Zaizov R., Varsano L. 1971. Postnatal changes in some red cell parameters. *Acta paediat Scand*, 60.

Mantovani, R., Malgaretti, N., Nicolis, S., Giglioni, B., Comi, P., Cappellini, N., Bertero, M.T., Caligaris-Cappio, F. & Ottolenghi, S. (1988). An erythroid specific nuclear factor binding to the proximal CACCC box of the β -globin gene promoter. *Nucleic Acids Res*, 16, 4299-313.

Michelson A.M., & Orkin, S. H. (1980). The 3' untranslated regions of the duplicated human α -globin genes are unexpectedly divergent. *Cell*, 22, 371-337.

Modell, C.B., & Berdoukas, V.A. (1984). The clinical approach to thalassemia. *Ann N Y Acad Sci*, 232, 201.

Proudfoot, N.J. (1979). Eukaryotic promoters. *Nature*, 279, 376.

Nadkarni, A., Gorakshakar, A.C., Lu, C.Y., Krishnamoorthy, R., Ghosh, K., Colah, R. & Mohanty, D. (2001). Molecular pathogenesis and clinical variability of β -thalassemia syndromes among Indians. *Am J Hematol*, 68, 75-80.

Nopparatana, C., Panich, V., Saechan, V., Srioongrueng, V., Nopparatana Cd., Rungjeadpha, J., Pornpatkul, M., Laosombat, V., & Fukumaki, Y. (1995). The spectrum of β -thalassemia mutations in southern Thailand. *Southeast Asian J Trop Med Public Health*, 26 (suppl. 1), 229-234.

Orkin S. H., & Michelson, A. (1980). Partial deletion of the globin structural gene in human α thalassemia. *Nature*, 286, 538.

Orkin, S.H., Kazazian, H.H., Jr., Antonarakis, S.E., Goff, S.C., Boehm, C.D., Sexton, J.P., Waber, P.G. & Giardina, P.J. (1982). Linkage of β -thalassaemia mutations and β -globin gene polymorphisms with DNA polymorphisms in human β -globin gene cluster. *Nature*, 296, 627-31.

Orkin, S.H., Sexton, J.P., Cheng, T.C., Goff, S.C., Giardina, P.J., Lee, J.I. & Kazazian, H.H., Jr. (1983). ATA box transcription mutation in β -thalassemia. *Nucleic Acids Res*, 11, 4727-34.

Poncz, M., Ballantine, M., Solowiejczyk, D., Barak, I., Schwartz, E. & Surrey, S. (1982). β -thalassemia in a Kurdish Jew. Single base changes in the TATA box. *J Biol Chem*, 257, 5994-6.

Pressley, L., Higg, D.R., Clegg, J.B., & Weatherall, D.J. (1980). Gene deletion in α -thalassemia prove that the 5' ζ locus is functional. *Proc Natl Acad Sci U S A*, 77, 3586.

Proudfoot N.J., & Brownlee, G.G. (1976). 3' non-coding sequences in eukaryotic messenger RNA. *Nature*, 263, 211-214.

Qatanani, M., Taher, A., Koussa, S., Naaman, R., Fisher, C., Rugless, M., Old, J. & Zahed, L. (2000). β -thalassaemia intermedia in Lebanon. *Eur J Haematol*, 64, 237-44.

Rund, D., Oron-Karni, V., Filon, D., Goldfarb, A., Rachmilewitz, E. & Oppenheim, A. (1997). Genetic analysis of β -thalassemia intermedia in Israel: diversity of mechanisms and unpredictability of phenotype. *Am J Hematol*, 54, 16-22.

Rust, S., H. Funke and G. Assmann. (1993). Mutagenically separated PCR (MS-PCR): a highly specific one step procedure for easy mutation detection. *Nucleic Acids Res*, 21, 3623-3629.

Sampietro, M., Thein, S.L., Contreras, M., & Pazmany, L. (1992). Variation of HbF and F-cell number with the $^G\gamma Xmn$ I (C-T) polymorphism in normal individuals. *Blood*, 79(3), 832-3.

Sanguansermsri, T., Thanarattanakorn, P., Steger, H.F., Tongsong, T., Chanprapaph, P., Wanpirak, C., Siriwanapan, P., Sirichotiyakul, S. & Flatz, G. (2001). Prenatal diagnosis of β -thalassemia major by high-performance liquid chromatography analysis of hemoglobins in fetal blood samples. *Hemoglobin*, 25, 19-27.

Sanguansermsri, T., Pape, M., Laig, M., Hundrieser, J., Pape M., & Flaz, G. (1990).

β^0 -thalassemia in Thai family is caused by 3.4 kb deletion including the entire β -gene.

Hemoglobin, 14, 157-168.

Sanguansermsri, T., Phumyu, N., Chomchven, S., & Steger, HF. (1999). Screening for α -thalassemia-1 heterozygotes in expecting couples by combination of a simple erythrocyte osmotic fragility test and a PCR-based method. *Community Genet*, 2, 26-39.

Schroeder, W.A., Huisman, T.H.J., Brown, A.K., Uy, R., Bouwer, N.G., Leerch, P.O., Shelton, J.R., Shelton, J.B., & Apell, G. (1971). Postnatal changes in the chemical heterogeneity in the fetal hemoglobin. *Pediatr Res*, 5, 493.

Sirichotiyakul, S., Saetung, R., Sanguansermsri, T. (2003). Analysis of β -thalassemia mutations in Northern Thailand using an automated fluorescence DNA sequencing technique. *Hemoglobin*, 27, 89-95.

Smith, EW., & Torbert, JB. (1958). Study of two abnormal hemoglobins with evidence for a new genetic locus for hemoglobin formation. *Bull Johns Hopkins Hosp.*, 102, 38-48.

Sturgeon, Itano, H.A., & Bergren W.R. (1955). Genetic and biochemical studies of "intermediate" types of Cooley's anemia. *Br J Haematol*, 1.

Takihara, Y., Forget, B.G., Scarpa, A., & Benz E.J. (1984). Intranuclear defect in β -globin mRNA accumulation due to a premature translation termination codon. *Blood*, 64, 13.

Thein S.L., & Reittie, J.E. (1998). F cells by immunofluorescent staining of erythrocyte smears. *Hemoglobin*, 22, 415-7.

Thein, S.L., Hesketh, C., Wallace, R.B. & Weatherall, D.J. (1988). The molecular basis of thalassaemia major and thalassaemia intermedia in Asian Indians: application to prenatal diagnosis. *Br J Haematol*, 70, 225-31.

Thein, S.L., Old, J.M., Wainscoat, J.S., Petrou, M., Modell, B. & Weatherall, D.J. (1984). Population and genetic studies suggest a single origin for the Indian deletion beta thalassaemia. *Br J Haematol*, 57, 271-8.

Traeger, J., Wood, W.G., Clegg, J.B., Weatherall ,D.J., & Wasi, P. (1980). Defective synthesis of Hb E is due to reduced level of β^E mRNA. *Nature*, 288, 497.

Treisman, R., Orkin, S.H., & Maniatis T. (1983). Specific transcription and RNA splicing defects in five cloned β -thalassemia genes. *Nature*, 302, 591-596.

Trent, RJ., Higgs, DR., Clegg JB., & Weatherall, DJ. (1981). A new triplicated α -globin gene arrangement in man. *Br J Haematol*, 49, 149-152.

Tuan, D., Solomon, W., Li, Q., & London, I.M. (1985). The & quote; β -like-globin & quot; gene domain in human erythroid cells. *Proc Natl Acad Sci U S A*, 82, 6384-8.

Wainscoat, J.S., Kanavakis E., Wood W.G., & et al. (1983). Thalassemia intermedia in Cyprus-the interaction of α - and β -thalassemia. *Br J Haematol*, 53, 411.

Walsh, PS., M.D., Higuchi R. (1991). Chelex-100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *Biotechniques*, 10, 506-13.

Weatherall, D.J. (1981). The Thalassemia Syndromes. Blackwell Scientific: Oxford.

Weatherall, D.J. (2001a). Phenotype-genotype relationships in monogenic disease: lessons from the thalassaemias. *Nat Rev Genet*, 2, 245-55.

Weatherall, D.J. (2001b). The Thalassemias. In *Williams Hematology (sixth edition)*, McGraw-Hill: New York.

Winichagoon, P., Echaroen, S., Chen, P. & Wasi, P. (2000). Genetic factors affecting clinical severity in β -thalassemia syndromes. *J Pediatr Hematol Oncol*, 22, 573-80.

Wintrobe, M. (1885,1981). *Clinical hematology*.: Philadelphia,USA.

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright[©] by Chiang Mai University
All rights reserved